

Hypoalbuminemia and nodular duodenal mucosa

Maddalena Zippi*; Sirio Fiorino; Innocenzo Bertoldi; Wandong Hong; Giuseppe Occhigrossi

*Maddalena Zippi, MD, PhD

Unit of Gastroenterology and Digestive Endoscopy, Sandro Pertini Hospital, Via dei Monti Tiburtini 385, 00157 Rome, Italy

Phone: +39-06-41433310, Fax: +39-06-41433847; Email: maddyzip@yahoo.it

Keywords

Common variable immunodeficiency (cvid); *giardia lamblia*; malabsorption; nodular lymphoid hyperplasia (nlh)

Clinical Image Description

A 26-year-old Egyptian man was admitted to the Emergency Department of the Sandro Pertini Hospital due to acute peripheral oedema associated with weight loss (6 kg in the last 7 months) and chronic diarrhoea. In the anamnesis, no previous episodes of infections (gastrointestinal or respiratory) or hospitalizations were reported. Physical examination of the patient revealed his poor condition. Hypoalbuminemia (2.6 g/dl) and anaemia (haemoglobin 10.5 g/dl) were present. Serological screening for celiac disease was negative. A gastroscopy and a colonoscopy with ileoscopy (Figure 1) with histological examinations were performed. Gastroscopy showed diffuse macroscopic nodular appearance on bulb and duodenum, while ileoscopy revealed the same nodular pattern but less represented. Histopathological examination of the duodenal biopsies displayed nodular lymphoid hyperplasia (NLH) associated with *Giardia lamblia* infection (Figure 2). This finding was confirmed by parasitic examination of the faeces, which highlighted the presence of *Giardia* cysts. Otherwise, *Helicobacter pylori* was not present in gastric biopsies. Crohn's disease was excluded both with the histological examination of the terminal ileum and owing to the negativity of the fecal calprotectin dosage. Further examination showed low levels of quantitative immunoglobulin (IgG 240 mg/dL, IgA 21 mg/dL, and IgM 36 mg/dL), mutation of the gene TNFRSF13B, and immunological consultation. According to all these evidences, a common variable immunodeficiency complicated by *Giardia lamblia* infection was diagnosed. The patient was treated with bowel rest, intravenous albumin, parenteral nutrition, as well as immunoglobulin replacement and antibiotic therapy (metronidazole).

Discussion

Common variable immunodeficiency (CVID) is a group of disorders with nonspecific clinical mani-

festations, largely relying on the decreased serum immunoglobulin levels [1]. Its epidemiology and its prevalence are very heterogeneous. The last one has been accounted for 40.2% in the USA, 2.6% in the Middle East countries and 1.3% in the Africa [1]. This condition manifests itself in the course of the second/third decade of life, affecting about 1:50.000-1:200.000 individuals in the general population, whose immune system is not able to produce antibodies against bacteria, viruses or other affecting the body [2]. Over 90% of CVID patients are more susceptible to contract bacterial infections, especially those affecting the respiratory and gastrointestinal tract (*Giardia lamblia* followed by *Salmonella* and *Campylobacter jejuni*) [2]. *Giardia lamblia* infection can generate a nodular mucosal pattern, which may look like a celiac disease or a Crohn disease when observed from an endoscopic point of view [3]. It has been reported that about half of untreated CVID patients suffer from malabsorption, and NLH may also be present [4]. This constitutes a risk factor for the development of intestinal lymphoma and entailing the implementation of endoscopic surveillance protocols [5].




Figure 1: Endoscopy showing the mucosa of : a) bulb; b) duodenum; c) terminal ileum.




Figure 2: Biopsy of the duodenal mucosa (haematoxylin & eosin 10X) showing: a) increase in lymphoid infiltrate in the lamina propria organized in follicles (white arrow); b) falciform corpuscles (White arrow) compatible with the presence of *Giardia*.

References

1. Modell V, Orange JS, Quinn J, Modell F. Global report on primary immunodeficiencies: 2018 update from the Jeffrey Modell Centers Network on disease classification, regional trends, treatment modalities, and physician reported outcomes. *Immunol Res* 2018; 66: 367-380.
2. Cunningham-Rundles C, Bodian C. Common variable immunodeficiency: clinical and immunological features of 248 patients. *Clin Immunol* 1999; 92: 34-48.
3. Ranchod M, Lewin KJ, Dorfman RF. Lymphoid hyperplasia of the gastrointestinal tract. A study of 26 cases and review of the literature. *Am J Surg Pathol.* 1978; 2: 383-400.
4. Khodadad A, Aghamohammadi A, Parvaneh N, et al. Gastrointestinal manifestations in patients with common variable immunodeficiency. *Dig Dis Sci.* 2007; 52: 2977-2983.
5. Abolhassani H, Sagvand BT, Shokuhfar T, et al. A review on guidelines for management and treatment of common variable immunodeficiency. *Expert Rev Clin Immunol.* 2013; 9: 561-575.

Manuscript Information: Received: September 15, 2019; Accepted: October 16, 2019; Published: October 31, 2019

Authors Information: Maddalena Zippi^{1*}; Sirio Fiorino²; Innocenzo Bertoldi³; Wandong Hong⁴; Giuseppe Occhigrossi¹

¹Unit of Gastroenterology and Digestive Endoscopy, Sandro Pertini Hospital, Rome, Italy

²Unit of Internal Medicine, Maggiore Hospital, Local Health Unit of Bologna, Bologna, Italy

³Unit of General Surgery, Sandro Pertini Hospital, Rome, Italy

⁴Department of Gastroenterology and Hepatology, First Affiliated Hospital of Wenzhou Medical University, Wenzhou City, Zhejiang, The People's Republic of China

Citation: Zippi M, Fiorino S, Bertoldi I, Hong W, Occhigrossi G. Hypoalbuminemia and nodular duodenal mucosa. *Open J Clin Med Case Rep.* 2019; 1594.

Copy right statement: Content published in the journal follows Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>). © **Zippi M 2019**

About the Journal: Open Journal of Clinical and Medical Case Reports is an international, open access, peer reviewed Journal focusing exclusively on case reports covering all areas of clinical & medical sciences.

Visit the journal website at www.jclinmedcasereports.com

For reprints and other information, contact info@jclinmedcasereports.com