

Arthroscopic management of articular hoffa fracture: A case report

Adnane Lachkar*; Anne Marie Abassa Kede; Hicham Yacoubi; Abdeljaouad Najib

***Corresponding Author: Adnane Lachkar**

Department of orthopedic surgery B, Hospital University Center Mohamed VI, University Mohamed first Oujda, Morocco.

Phone: (212) 0669021976 / (33) 0644000830; Email: dr.lachkar@gmail.com

Abstract

Hoffa fractures are uncommon. These joint fractures require anatomical reduction with stable osteosynthesis allowing early rehabilitation. The arthroscopic reduction with percutaneous osteosynthesis has the advantage of being a minimally invasive technique allowing a good control of the reduction with minimal morbidity. We report the case of a patient treated for Hoffa's fracture by percutaneous fixation arthroscopically assisted with a satisfying outcome.

Keywords

Hoffa fracture; knee articular fractures; condyle fractures; arthroscopy

Introduction

Fractures of the distal femur are a rare occurrence; they represent less than 10% of all femoral fracture. Unicondylar fractures represent approximately 3% and Hoffa fractures are an even unusual entity; challenging in terms of management. These joint fractures require anatomical reduction with stable osteosynthesis allowing early rehabilitation [1,2]. The arthroscopic reduction with percutaneous osteosynthesis of Hoffa fractures has the advantages of being a minimally invasive technique with minimal morbidity. It also offers a good control of the reduction and allows an appropriate treatment at the same time [3,4]. In recent literature, the published series concerning arthroscopic management of this type of fracture remain limited. We report the case of a patient treated for Hoffa's fracture by percutaneous osteosynthesis under arthroscopic control.

Case Report

A 35-year-old patient with a history of left knee traumatic injury after a motorbike accident. His

Clinical examination showed a painful and swollen knee (Figure 1). The initial imaging evaluation consisting of anteroposterior and a lateral plain radiographs revealed a displaced Hoffa fracture of the lateral femoral condyle. Computed tomography scan confirmed the diagnosis (Figure 2).



Figure 1: X-rays; Hoffa fracture of lateral condyle.



Figure 2: Computed tomography scan; Hoffa fracture of lateral condyle .

After due informed consent, patient was planned for arthroscopic-assisted fixation of his fracture. In the operating room, he was positioned supine with a tourniquet to the upper thigh. After evacuation of the hemarthrosis, diagnostic arthroscopy was envisaged (Figure 3). The menisci and cruciate ligaments were found to be regular. Joint exploration showed a large Hoffa fracture with osteochondral lesions. The clots were removed by a 3.5 mm shaver and loose fragments were excised (Figure 4).




Figure 3: A- Evacuation of the hemarthrosis. B- Diagnostic arthroscopy.




Figure 4: Removing the clots between the fracture surfaces by a 3.5 mm shaver.

Fracture reduction was achieved with K-wires used as a joystick. Then, two guide wires were placed in anteroposterior direction (perpendicular to fracture site). The length and good position of the Kirschner wires were verified under C-arm. Osteosynthesis was carried out by 6.5 mm cannulated screws. Osteochondral lesions were repaired with Herbert screws (Figures 5,6). The reduction was anatomical and the osteosynthesis was stable (Figure 7).

The Postoperative management was simple. Rehabilitation was immediately started with a good recovery of the articular range of motion. The Patient was allowed partial weight bearing at 6 week and total weight bearing was allowed by the 3rd month. Functional results were very satisfactory.




Figure 5: Osteosynthesis of the fracture by 6.5mm cannulated screws




Figure 6: Reparation of osteochondral lesions with Herbert screws




Figure 7: Final fluoroscopic control

Discussion

In 1904, Albert Hoffa was the first to describe the typical form for fractures of the femoral condyle in the sagittal plane [5]. The therapeutic management of these injuries has undergone a large evolution over the years. Indeed, reduction techniques with internal fixation assisted by arthroscopy appear as new methods which provide several advantages [3,4]:

- Surgical aggression is relatively minimal, thus avoiding damage to surrounding tissue, including the risk of iatrogenic damage to the common peroneal nerve.
- The joint architecture can be inspected thoroughly by arthroscopic control. Consequently, associated lesions (in particular cartilage or ligament damages) can be treated simultaneously.
- The use of a multi-angle screw fixation under fluoroscopic control offers a stable and reliable fixation preventing any risk of rotation or second displacement of the fragment.
- The operating time is shorter, the bleeding is minimal and the risk of infection is so negligible because the procedure is performed under continuous irrigation.
- The scar is minimal and conducive to the rapid recovery of the knee joint range of motion.

The procedure is performed under spinal anaesthesia; patient is installed in supine position with a tourniquet to the upper thigh. Conventional arthroscopic approaches are used. After evacuating the hemarthrosis, a first diagnostic-look is made. The cartilage, menisci and ligaments lesions are checked. The clots between the fracture surfaces are excised by a shaver.

Generally, the condylar fragment moves retro-femur due to its posterior capsular attachments. A 120° knee flexion relaxes the posterior capsule and the gastrocnemius muscles, thereby reducing the fracture using a simple curved Halsted forceps. A Kirschner wire can be used as a joystick to assist reduction

manoeuvres. Also, the tibial plateau contributes to the repositioning of the fragment by pushing it back during hyperflexion. The quality of reduction is carefully controlled by the magnification offered by the arthroscope camera. The condyle is then temporarily stabilized with two guide pins inserted obliquely and oriented medially and anteriorly from the most posterolateral point of the fragment, just above the lateral meniscus. The final fixation is achieved by 6.5 mm cannulated screws following the path of the guide wires. A minimum of 2 screws is required to ensure the rotational stability of the fragment. The screw heads must be inserted through the articular cartilage. The small comminuted fragments are generally removed while the larger are anatomically fixed using Herbert screws under arthroscopic control. The reduction and alignment of the screws are confirmed by intraoperative fluoroscopy. A stability test is then performed by bending the knee up to 140° [3,4].

In the postoperative, walking without bear weight is allowed on the second day of surgery. Full weight bearing is allowed from the 12th week. The functional results are globally satisfactory and remarkable [3,4,6] which makes this surgical technique as an excellent therapeutic option in correctly selected patients (Table 1).

Indications	- Recent fractures: mono-fragment, large volume - Neglected fractures: without comminution, large fragment with good bone quality
Contraindications	- Recent fractures: Irreducibility or comminution - Neglected fractures: Pseudarthrosis requiring grafting

Few reports of arthroscopic assisted fixation of Hoffa fractures are available [7–11]. Wallenbock and Ledinski [7] reported on a series of 24 intraarticular knee fractures surgically treated two Hoffa injuries. They confess the demanding and challenging nature of the arthroscopic fixation. However, they report good early results. McCarthy [8] reported a sagittal plane lateral femoral condyle fracture. They were not able to achieve adequate reduction but acclaimed the decreased blood loss, soft tissue dissection, operative time, and the accelerated recovery time.

Arthroscopy has been described as a useful tool in the treatment of neglected Hoffa fractures without bone loss. It also wells as a great treatment for intra-articular and ligamentous injuries which could be not recognized by clinical exam, computed tomography scan or magnetic resonance imaging [1,4,10].

Conclusion

Arthroscopic-assisted osteosynthesis of Hoffa fractures is a technically demanding surgical method, most often applicable for fresh and simple fractures. However, despite its long learning curve, its limited and poorly coded indications, this treatment promises multiple advantages. It reduces the rate of complications encountered with open surgery and thus offering broad prospects for the management of these joint fractures.

References

1. Zhou Y, Pan Y, Wang Q, Hou Z, Chen W. Hoffa fracture of the femoral condyle: Injury mechanism, classification, diagnosis, and treatment. *Medicine (Baltimore)*. 2019; 98(8): e14633.
2. Patel PB, Tejwani NC. The Hoffa fracture: Coronal fracture of the femoral condyle a review of literature. *J Orthop*. 2018; 15(2): 726-731.
3. Xiao K, Chen C, Yang J, Yang D, Liu J. An attempt to treat Hoffa fractures under arthroscopy: A case report. *Chin J Traumatol*. 2018; 21(5): 308-310.
4. Goel A, Sabat D, Agrawal P. Arthroscopic-assisted fixation of Hoffa fracture: A case report and description of technique. *J ClinOrthop Trauma*. 2016; 7(1): 61-5.
5. Stuttgart Hoffa Afa. *Lehrbuch der Frakturen und Luxationen*. Verlag von Ferdinand Enke; 1904: 451.
6. Say F, Kuyubaş N. Arthroscopy-Assisted Fixation of an Intra-Articular Medial Femoral Condyle (Hoffa) Fracture of the Knee: A Case Report. *JBJS Case Connect*. 2014; 4(4): e109.
7. Wallenbock E, Ledinski C. Indications and limits of arthroscopic management of intraarticular fractures of the knee joint. *Aktuelle Traumatologie*. 1993; 23:97-101 [in German].
8. McCarthy JJ, Parker RD. Arthroscopic reduction and internal fixation of a displaced intraarticular lateral femoral condyle fracture of the knee. *Arthroscopy*. 1996; 12: 224-227.
9. Stern RE. Comment. *Arthroscopy*. 1996; 12: 760-761.
10. Demirel M, Dereboy F, Ozturk A. Arthroscopically assisted intra-articular lateral femoral condyle fracture surgery. *Arthroscopy*. 2006; 22: 690e1-690e4 .
11. Ercin E, Bilgili MG, Basaran SH, et al. Arthroscopic treatment of medial femoral condylar coronal fractures and nonunions. *Arthrosc Tech*. 2013; 2(4): e413-e415.

Manuscript Information: Received: January 28, 2020; Accepted: April 24, 2020; Published: April 30, 2020

Authors Information: Adnane Lachkar*; Anne Marie Abassa Kede; Hicham Yacoubi; Abdeljaouad Najib
Department of orthopedic surgery B, Hospital University Center Mohamed VI, University Mohamed first Oujda, Morocco.

Citation: Lachkar A, Abassa Kede AM, Yacoubi H, Najib A. Arthroscopic management of articular Hoffa fracture: A case report. *Open J Clin Med Case Rep*. 2020; 1656.

Copy right statement: Content published in the journal follows Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>). © **Lachkar A 2020**

About the Journal: Open Journal of Clinical and Medical Case Reports is an international, open access, peer reviewed Journal focusing exclusively on case reports covering all areas of clinical & medical sciences.

Visit the journal website at www.jclinmedcasereports.com

For reprints and other information, contact info@jclinmedcasereports.com